

IFP-465-UNC

hep-ph/9308275

August 1993

Cubic Mass Relation in $SU(3) \times U(1)$ Electroweak Theory

Paul H. Frampton, Plamen I. Krastev* and James T. Liu

Institute of Field Physics, Department of Physics and Astronomy,

University of North Carolina, Chapel Hill, NC 27599-3255, USA

Abstract

In the 331 model, lepton number may be explicitly broken by trilinear scalar self couplings. This leads to neutrino masses proportional to the cube of the corresponding charged lepton mass, with consequences for solar neutrinos and for hot dark matter.

*Permanent address: Institute of Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, BG-1784 Sofia, Bulgaria.

Whether neutrinos have non-zero rest mass is still an unresolved question. It is possibly the question whose answer will provide the first evidence of physics beyond the minimal standard model of particle interactions. Theoretically, there is no known symmetry reason for vanishing neutrino masses; experimentally, there are indications of non-vanishing neutrino masses which await confirmation. In the present letter we anticipate the emergence of non-zero neutrino masses and derive a cubic relation between neutrino and charged lepton masses.

Concerning the experimental situation, solar neutrino experiments [1] suggest a deficit of solar neutrinos compared to the predictions of the standard solar model; if this is interpreted as due to the Mikheyev–Smirnov–Wolfenstein (MSW) mechanism [2] one is led to expect $m(\nu_\mu) \sim 10^{-3}\text{eV}$ while $m(\nu_e)$ is much smaller [3]. Also, if ν_τ is a significant component of hot dark matter in the universe one expects [4] $m(\nu_\tau) \simeq 30\text{eV}$. Thirdly, the measurements of atmospheric ν_μ and ν_e neutrinos suggest an anomalously small muon neutrino flux which may be interpreted as due to non-zero neutrino masses of either ν_μ or ν_τ of the order of 10^{-1}eV [5].

These provocative hints from experiment lead inevitably to the question of how neutrino masses are generated in the theory. A popular conjecture is the see-saw mechanism [6] which naturally accounts for why neutrino masses are small. In a *quadratic* see-saw, $m(\nu_i)$ is proportional to $m(f)^2/M$ where f is some charged fermion in the same family and M is a larger mass scale; this is based on a grand unification scenario with a desert hypothesis and on the assumption that right-handed neutrino fields exist, one for each known fermion family. It is usually assumed that f is the up-quark member of the same family, based on $SO(10)$ arguments (see *e.g.* [7] and references therein).

Here we examine the question of neutrino masses in the 331 model [8,9]. In the minimal version of the 331 model there are no right-handed neutrinos. Furthermore, while lepton family number is not conserved, total lepton number $L = L_e + L_\mu + L_\tau$ is conserved in the minimal 331 model. As a result, neutrinos remain massless just as in the minimal standard electroweak model.

It is straightforward to give neutrinos a Dirac mass in the 331 model by the introduction of right-handed neutrinos. However this approach has the same drawback as in the standard model, namely the Yukawa couplings have to be unnaturally small in order to account for the small neutrino masses. We are thus led to consider the case where neutrinos pick up a Majorana mass, which necessarily involves violating lepton number. Spontaneous breaking of L will lead to a massless triplet Majoron [10] which has been ruled out experimentally by Z lineshape measurements as mentioned briefly in [8]. Thus if L is broken it must be done explicitly.

We first review the relevant features of the 331 model. The electroweak gauge group is $SU(3)_L \times U(1)_X$ and the leptons are in antitriplets with $X = 0$, $L_{i\alpha} = (\ell_i^-, \nu_i, \ell_i^+)$ for $i = e, \mu, \tau$. The quark sector is unimportant for the following discussion, but, via anomaly cancellation, gives rise to the three family requirement. Symmetry breaking and fermion masses arise from four Higgs multiplets, of which only two couple to the leptons. These are a sextet Higgs $H^{\alpha\beta}$ and a triplet Higgs ϕ^α both with $X = 0$. When $SU(3)_L \times U(1)_X$ is broken down to $SU(2)_L \times U(1)_Y$, the sextet decomposes as $\mathbf{6}_0 \rightarrow \mathbf{3}_2 + \mathbf{2}_{-1} + \mathbf{1}_{-4}$ into $SU(2)$ triplet, doublet and singlet Higgs, and the triplet decomposes as $\mathbf{3}_0 \rightarrow \mathbf{2}_1 + \mathbf{1}_{-2}$ into a doublet and singlet.

The lepton Yukawa couplings may be written

$$- \mathcal{L} = h_1^{ij} L_\alpha^i L_\beta^j H^{\alpha\beta} + h_2^{ij} L_\alpha^i L_\beta^j \bar{\phi}_\gamma \epsilon^{\alpha\beta\gamma} + \text{h.c.} , \quad (1)$$

where the primes denote weak eigenstates. Here h_1^{ij} is symmetric in family space while h_2^{ij} is antisymmetric. In order for the Yukawas to conserve L , the $SU(2)$ triplet and singlet Higgs must carry lepton numbers -2 and 2 respectively while the doublets have $L = 0$.

In the minimal 331 model, charged lepton masses arise from the scalar VEVs $\langle \phi^2 \rangle = v/\sqrt{2}$ and $\langle H^{13} \rangle \equiv \langle H^{31} \rangle = y/2$. The resulting mass matrix is given in family space by

$$m(\ell)^{ij} = h_1^{(ij)} y + \sqrt{2} h_2^{[ij]} v . \quad (2)$$

Assuming h_1^{ij} and h_2^{ij} are real so there is no CP violation in the leptonic sector, we may pick a basis where $h_1^{ij} = h_1^i \delta^{ij}$ is diagonal and h_2 remains antisymmetric. There are then a total

of six Yukawa parameters: three charged lepton masses and three mixing angles. To avoid unsuppressed lepton flavor violating processes [11], we demand $h_2 v \ll h_1 y$. In this case, we find charged lepton masses $m(\ell_i) \approx h_1^i y$ and mixing angles $\alpha^{ij} \approx \sqrt{2} h_2^{ij} v / (h_1^i + h_1^j) y$. In the minimal case where neutrinos remain massless, they can be rotated so the $W^\pm$ charged current is lepton family diagonal. However, the α^{ij} angles show up in the couplings to both singly and doubly charged dileptons [12].

We now give neutrinos masses by breaking L explicitly [13]. This is done in the Higgs sector by adding the two soft (dimension three) cubic self couplings

$$- \mathcal{L} = m_1 H^{\alpha_1 \beta_1} H^{\alpha_2 \beta_2} H^{\alpha_3 \beta_3} \epsilon_{\alpha_1 \alpha_2 \alpha_3} \epsilon_{\beta_1 \beta_2 \beta_3} + m_2 H^{\alpha \beta} \overline{\phi}_\alpha \overline{\phi}_\beta + \text{h.c.} . \quad (3)$$

Both terms violate L by two units. The cubic couplings in general induce a linear term in H^{22} and shifts the vacuum value to $\langle H^{22} \rangle = (3m_1 y^2 - m_2 v^2) / 2M(H^{22})^2$. This triplet Higgs VEV is constrained by the ρ -parameter, but, if present, gives rise to tree-level lepton number violation and Majorana neutrino masses. Since lepton number has been broken explicitly, there would be no massless Majoron in this case.

An interesting case occurs when $3m_1 y^2 = m_2 v^2$, which implies that at tree level the vacuum value of H^{22} vanishes. Imposing this requirement that $\langle H^{22} \rangle = 0$ prevents large tree-level lepton number violating processes and furthermore gives agreement with the experimentally observed limits on the ρ -parameter. Here we shall further assume that $\langle H^{22} \rangle$ remains zero even with radiative corrections. This necessarily involves fine tuning order by order in perturbation theory. This assumption is made to obtain an interesting new pattern of the neutrino masses which might be derived more naturally in some future different model.

In the minimal 331 model, with $m_1 = m_2 = 0$ in Eq. (3), lepton number remains an exact symmetry of the theory. In this case, with the assumptions we have made (especially $\langle H^{22} \rangle = 0$) the neutrinos cannot acquire mass. If we further set the antisymmetric Yukawa coupling h_2^{ij} to zero, then the Yukawa interactions can be taken to be family diagonal. As a result, only dileptons can violate lepton family number, and hence there is the selection

rule $\Delta L_i = 0 \bmod 2$.

Next consider turning on the soft lepton number violating terms, $3m_1 y^2 = m_2 v^2 \neq 0$ in Eq. (3), and keeping $h_2^{ij} = 0$ in Eq. (1). In this case, the lepton Yukawas remain family diagonal, and we may work in the basis for the L_α^i fields in which h_1^{ij} is diagonal, $h_1^{ij} = h_1^i \delta^{ij}$. Since we have imposed $\langle H^{22} \rangle = 0$, the neutrinos are massless at tree level. At one-loop level, however, the diagram of Fig. 1(a) gives a finite contribution to the neutrino masses.

Treating the lepton number violating mass insertion on the Higgs line, $\delta m^2 = 6m_1 y$, as a perturbation, the one-loop diagram is evaluated as

$$\begin{aligned} i\Sigma^i(p) &= 2 \int \frac{d^4 k}{(2\pi)^4} (i2h_1^i) \gamma_L \frac{i}{\not{k} + \not{p} - m_i} (i2h_1^i) \gamma_L \frac{i}{k^2 - M(H^{23})^2} (i\delta m^2) \frac{i}{k^2 - M(H^{12})^2} \\ &= -8(h_1^i)^2 m_i \delta m^2 \gamma_L \int \frac{d^4 k}{(2\pi)^4} \frac{1}{(k+p)^2 - m_i^2} \frac{1}{k^2 - M(H^{23})^2} \frac{1}{k^2 - M(H^{12})^2} , \end{aligned} \quad (4)$$

where i labels the lepton family and $\gamma_L = \frac{1}{2}(1 - \gamma^5)$ projects onto left-handed chiralities. The factor of 2 takes into account both the diagram and the conjugate one. It is straightforward to perform the loop integration in the limit $p \ll m_i \ll M(H)$. Using $m(\ell_i) = h_1^i y$, we arrive at the result

$$m(\nu_i) = C m(\ell_i)^3 / M_W^2 , \quad (5)$$

where the dimensionless constant C is given by

$$C = \frac{\alpha_2}{2\pi \sin^2 \beta} \frac{\delta m^2}{M(H^{23})^2 - M(H^{12})^2} \ln \left(\frac{M(H^{23})^2}{M(H^{12})^2} \right) . \quad (6)$$

We have defined $\tan \beta = y / \sqrt{v^2 + v'^2}$ where v , v' and y are the VEVs of the three Higgs doublets giving mass to the $W^\pm$, $M_W = \frac{g}{2} \sqrt{v^2 + v'^2 + y^2}$.

Eq. (5) is the cubic relationship between neutrino mass and charged lepton mass. Unlike the model of Ref. [14] where the various Yukawa couplings are unrelated, in the 331 model, the *same* Yukawa couplings, Eq. (1), are responsible for both the charged lepton mass and, through the loop diagram, the neutrino mass. This is the origin of the above cubic mass relationship. More generally, this feature of the 331 model leads to definite structures of both neutrino masses and mixing.

Family mixing in the 331 model is complicated both by the fact that there are additional couplings to dileptons and by the large number of scalars in the model. In order to simplify the discussion, we work with small mixing in the leptonic sector. In this case, we can parametrize the charged lepton mixing by the three angles α^{ij} as before. Diagonalizing the neutrino mass matrix gives rise to neutrino mixing angles θ^{ij} .

While neutral currents are always family diagonal in the leptonic sector, mixing angles in the charged currents are given approximately by $\alpha^{ij} - \theta^{ij}$ for the coupling to W^- , $\alpha^{ij} + \theta^{ij}$ for Y^- and $2\alpha^{ij}$ for Y^{--} (all in the limit of small mixing) [12]. The strongest constraints on α^{ij} come from the lack of observation of lepton family violating decays such as $\mu \rightarrow 3e$ whose branching ratio is below 10^{-12} [15]. This process may occur via tree-level Y^{--} exchange in the 331 model. Since the dilepton mass is bounded to be under a TeV [8,16], this implies $\alpha^{12} < 10^{-5}$, which means that, in Eq. (1), $h_2^{12}v < 10^{-5}h_1^2y$. Somewhat weaker limits can also be placed on first-third and second-third family mixing.

Since we have assumed $h_2^{ij} = 0$ in deriving the cubic relation above, only the first term of Eq. (1) is present and the Yukawa couplings can be made family diagonal. As a result, $\alpha^{ij} = \theta^{ij} = 0$ and there is *no mixing* of neutrino or charged lepton flavors. In order to obtain mixing and $\Delta L_i = \pm 1$ lepton family violation we must make $h_2^{ij} \neq 0$ which gives rise to non-diagonal mass matrices.

In this more general case, the neutrino mixing angles θ^{ij} are not independent, but are related to the α^{ij} because of the common Yukawa couplings. When $h_2^{ij} \neq 0$ the remaining graphs (b) through (d) depicted in Fig. 1 also contribute to the neutrino mass matrix. In general, the relation between the mixing angles depends on the ratio of the soft breaking masses m_1 and m_2 in Eq. (3) as well as the various Higgs masses. For $\alpha^{ij} \ll 1$, we find

$$\theta^{ij} \approx \left(\frac{m_2}{12m_1} \right) \alpha^{ij} , \quad (7)$$

with the exact relation depending on the complete Higgs spectrum. From Eq. (7) it follows that $\theta^{12} \sim 10^{-2}$ as demanded by the MSW solution is compatible with $\alpha^{12} < 10^{-5}$ if $m_2 > 10^4 m_1$. This inequality, together with the relationship between $m_1 y^2$ and $m_2 v^2$, implies

that $v < 10\text{GeV}$ and suggests that the top quark acquires most of its mass by a mechanism different from the standard Yukawa couplings of the electroweak model. This was hinted at in [8] but the details of the mass generation mechanism are still under investigation.

We now address the phenomenology of the cubic mass relation, Eq. (5). Since the constant C given in Eq. (6) depends on the soft breaking terms and can be made arbitrarily small, the cubic formula does not give any prediction for the absolute masses. Nevertheless, it gives definite predictions for the ratios of neutrino masses. Because of the cubic relation, the strongest bound on C comes from the ν_τ mass bound, $m(\nu_\tau) < 35\text{MeV}$. Non-zero mixing angles θ^{ij} will perturb the exact cubic relation, but the deviations are small for small mixing.

For $m(\nu_\tau)$ we may adopt the estimate $m(\nu_\tau) = 29.3\text{eV}$ from the hot dark matter model of Sciama [17]. The massive ν_τ is not stable, and will decay radiatively, *e.g.* by $\nu_\tau \rightarrow \nu_\mu \gamma$. Since the gauge boson contribution at one-loop is GIM suppressed [18], the dominant decay mode is via the scalar exchange diagrams of Fig. 1 with a photon line attached in all possible ways [19]. These one-loop diagrams give rise to a $(\nu_i \leftrightarrow \nu_j)$ transition dipole moment

$$\begin{aligned}\mu_{ij} &\approx \frac{e\alpha_2}{\pi \sin^2 \beta} \frac{\delta m^2}{M_W^2} \frac{m(\ell_i)^3}{M(H)^4} \theta^{ij} f(m(\ell_i), M(H)) \\ &\approx 2eC \frac{m(\ell_i)^3}{M_W^2 M(H)^2} \theta^{ij} f(m(\ell_i), M(H)) \\ &\approx 2e \frac{m(\nu_i)}{M(H)^2} \theta^{ij} f(m(\ell_i), M(H)) ,\end{aligned}\tag{8}$$

where we have used the cubic mass relation, Eq. (5), to arrive at the last line. This shows the intimate relation between the radiatively generated Majorana mass and the transition moment. In the above, we have taken L_i to be the heavier family and dropped terms of order $m(\ell_j)/m(\ell_i)$. $M(H)$ is a typical Higgs mass, and the function f is given by

$$f(m(\ell_i), M(H)) \approx 2 + \ln \left(\frac{m(\ell_i)^2}{M(H)^2} \right) .\tag{9}$$

Note that this gives a moderate enhancement to the transition moment since typically $m(\ell) \ll M(H)$.

Assuming CP invariance, the transition dipole moment is either electric or magnetic, depending on the CP eigenvalues of the neutrinos [20]. It gives rise to a decay width

$$\Gamma(\nu_i \rightarrow \nu_j \gamma) = \frac{1}{8\pi} m(\nu_i)^3 |\mu_{ij}|^2, \quad (10)$$

assuming $m(\nu_i) \gg m(\nu_j)$. For ν_τ , the resulting radiative decay lifetime is given by

$$\tau \approx \frac{1}{2\alpha} \frac{M(H)^4}{m(\nu_\tau)^5} \frac{1}{(\theta^{23})^2} \frac{1}{f^2(m(\tau), M(H))}, \quad (11)$$

giving lifetimes as short as $\sim 10^{23}$ s as required by the cosmological scenario of Ref. [17]. Using the charged lepton masses then gives from Eq. (5) $m(\nu_\mu) = 6.2 \times 10^{-3}$ eV and $m(\nu_e) = 6.9 \times 10^{-10}$ eV. These neutrino masses are well below current experimental limits and are nicely consistent with the MSW mechanism [2].

This large scalar enhancement to the ν_τ decay rate also occurs in the charged leptonic sector for radiative processes such as $\mu \rightarrow e\gamma$. Although this is strongly constrained, the smallness of $m(\nu_\mu)$ from the cubic formula gives a predicted branching ratio well under current experimental limits. Thus bounds on the tree level dilepton contribution to $\mu \rightarrow 3e$ remain the strongest limit on lepton family mixing in the 331 model.

The effective Majorana mass of ν_e , $\langle m(\nu_e) \rangle = \sum_i m(\nu_i)(\delta^{1i} + \theta^{1i})^2$, leads to a contribution to neutrinoless double beta decay, $(\beta\beta)_{0\nu}$. Assuming $\theta^{13} < 10^{-2}$ since only $m(\nu_\tau)$ is significant, this contribution is at least three orders of magnitude below the current experimental limit [21]. In the 331 model, there are additional contributions to $(\beta\beta)_{0\nu}$ such as those indicated in the diagrams of Fig. 2. We estimate, however, that these dilepton diagrams are also many orders of magnitude beyond current empirical limits.

We have seen that the minimal 331 model is easily extended to give a highly predictive framework for generating neutrino masses and mixing. The basis for this lies in the fact that all leptons in a single family fall in a single SU(3) triplet with mass generation coming from common Yukawa couplings. Our principal result is the cubic mass relation, Eq. (5). It provides a consistent relation between $m(\nu_\tau)$ as required for hot dark matter and $m(\nu_\mu)$ as needed for the solar neutrino deficit. It further predicts $m(\nu_e)$ of about 0.7×10^{-9} eV.

The derivation we have given is flawed only by our treatment of the SU(2) triplet Higgs. A fine tuning was necessary to prevent the neutral component, H^{22} , from picking up a VEV

and hence giving the neutrinos mass at tree-level. If $\langle H^{22} \rangle$ is non-vanishing, the tree-level neutrino masses are of the form $m(\nu_i) \approx m(\ell_i) \delta m^2 / 4M (H^{22})^2$, assuming the triplet VEV takes on its natural scale. This then dominates the cubic term on the right hand side of Eq. (5) and leads to a linear mass relation which allows for a simultaneous solution in terms of neutrino oscillations of both the solar and atmospheric neutrino puzzles. Nevertheless, finding a model like [8,9] but where the neutral component of the triplet scalar is either absent or has a vacuum value constrained to vanish by a symmetry might be a fruitful line to pursue to make a rigorous derivation of a cubic mass relation.

We finally mention that a cubic mass relation can be accommodated by the see-saw mechanism using an inverse mass hierarchy for the heavy singlet Majorana masses [4]. However, the present cubic mass relation, Eq. (5), is not generated by a see-saw mechanism and does not involve right-handed neutrino fields. This relation comes instead because the 331 model provides a common framework for both charged leptons and neutrinos, once again showing the highly predictive aspects of this theory.

This work was supported in part by the U.S. Department of Energy under Grant No. DE-FG05-85ER-40219.

REFERENCES

- [1] R. Davis Jr., *et al.*, XXI Int. Cosmic Ray Conf., January 6–19, Adelaide, Australia, Conference papers, vol. 7, p. 155 (1990); Y. Totsuka, Talk given at the “Neutrino ’92” Int. Conf., Granada, June 8–13, 1992; A. I. Abazov *et al.*, (SAGE collaboration), Phys. Rev. Lett. **67** (1991) 332, and V. Gavrin, Talk given at the XXVIth Int. Conference on High Energy Physics, August 5–13, 1992, Dallas (U.S.A.); P. Anselmann *et al.*, (GALLEX collaboration), Phys. Lett. B **285** (1992) 376; **285** (1992) 390.
- [2] L. Wolfenstein, Phys. Rev. D **17** (1978) 2369; S. P. Mikheyev and A. Yu. Smirnov, Yad. Fiz. **42** (1985) 1441.
- [3] S. A. Bludman, N. Hata, D. C. Kennedy and P. G. Langacker, Phys. Rev. D **47** (1993) 2220; L. Krauss, E. Gates and M. White, Phys. Lett. B **299** (1993) 94; P. Krastev and S. T. Petcov, Phys. Lett. B **299** (1993) 99; X. Shi, D. N. Schramm and J. Bahcall, Phys. Rev. Lett. **69** (1992) 717.
- [4] H. Harari and Y. Nir, Nucl. Phys. B **292** (1987) 251.
- [5] K. S. Hirata *et al.*, Phys. Lett. B **280** (1992) 146; D. Casper *et al.*, Phys. Rev. Lett. **66** (1991) 2561; M. Goodman, ANL-HEP-CP-93-33.
- [6] H. Fritzsch and P. Minkowski, Phys. Lett. **62B** (1976) 72; M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, D. Z. Freeman and P. van Nieuwenhuisen, eds., North Holland, Amsterdam (1979); T. Yanagida, Prog. Th. Phys. B **135** (1978) 66.
- [7] S. A. Bludman, D. C. Kennedy and P. G. Langacker, Phys. Rev. D **45** (1992) 1810.
- [8] P. H. Frampton, Phys. Rev. Lett. **69** (1992) 2889.
- [9] F. Pisano and V. Pleitez, Phys. Rev. D **46** (1992) 410.
- [10] G. B. Gelmini and M. Roncadelli, Phys. Lett. **99B** (1981) 411.
- [11] S. Glashow and S. Weinberg, Phys. Rev. D **15** (1977) 1958.

- [12] J. T. Liu, to be published.
- [13] V. Pleitez and M. D. Tonasse, *Neutrinoless double beta decay in an $SU(3)_L \otimes U(1)_N$ model*, preprint hep-ph/9302201 (February 1993).
- [14] A. Zee, Phys. Lett. **93B** (1980) 389.
- [15] Particle Data Group, Phys. Rev. D **45** (1992) VI.15.
- [16] D. Ng, *The Electroweak Theory of $SU(3) \times U(1)$* , TRIUMF preprint TRI-PP-92-125 (December 1992).
- [17] D. W. Sciama, Ap. J. **364** (1990) 549; Int. J. of Mod. Phys. D **1** (1992) 161; SISSA preprint 34/93/A (February 1993) to appear in Ap. J. Lett..
- [18] W. J. Marciano and A. I. Sanda, Phys. Lett. **67B** (1977) 303; S. T. Petcov, Sov. J. Nucl. Phys. **25** (1977) 340; E: **25** (1977) 698.
- [19] S. T. Petcov, Phys. Lett. **115B** (1982) 401.
- [20] B. Kayser, Phys. Rev. D **30** (1984) 1023; P. B. Pal and L. Wolfenstein, Phys. Rev. D **25** (1982) 766.
- [21] W. C. Haxton, Talk given at the “Neutrino ’92” Int. Conf., Granada, June 8–13, 1992.

FIGURES

FIG. 1. One-loop diagrams contributing to neutrino mass. In addition, there are four more diagrams with the charged lepton replaced by its charge conjugate.

FIG. 2. Some of the potentially most significant additional contributions to neutrinoless double beta decay in the 331 model.